

# **Large-Class Strategies**

David E. Meltzer

Department of Physics and Astronomy  
ISU

# Outline

- Motivation and description of active-engagement teaching strategy.
- Watch video (18 minutes): pauses for comments and questions
- Describe details of questioning strategies
- Group work: Work with others in your (or related) discipline to create question sequences.

# Research in physics education and other scientific and technical fields suggests that:

- “Teaching by telling” has only limited effectiveness
  - listening and note-taking have relatively little impact
- Problem-solving activities with rapid feedback yield improved learning gains
  - student group work
  - frequent question-and-answer exchanges with instructor

***Goal: Guide students to “figure things out for themselves” as much as possible***

# What needs to go on in class?

- Clear and organized presentation by instructor is not *at all* sufficient
- Must find ways to guide students to synthesize concepts in their own minds
- Instructor's role becomes that of guiding students to ask and answer useful questions
  - aid students to work their way through complex chains of thought

# The Biggest Challenge: Large Lecture Classes

- Very difficult to sustain active learning in large classroom environments
- Two-way communication between students and instructor becomes paramount obstacle
- Curriculum development must be matched to innovative instructional methods

# Active Learning in Large Physics Classes

- **De-emphasis of lecturing**; Instead, ask students to respond to many questions.
- Use of classroom communication systems to obtain **instantaneous feedback** from entire class.
- Cooperative **group work** using carefully structured free-response worksheets

**Goal:** *Transform large-class learning environment into “office” learning environment (i.e., instructor + one or two students)*

# “Fully Interactive” Physics Lecture

*DEM and K. Manivannan, Am. J. Phys. 70, 639 (2002)*

- Very high levels of student-student and student-instructor interaction
- Simulate one-on-one dialogue of instructor’s office
- Use numerous structured question sequences, focused on specific concept: small conceptual “step size”
- Use student response system to obtain instantaneous responses from all students simultaneously (e.g., “flash cards”)





# Features of the Interactive Lecture

- High frequency of questioning
- Must often create unscripted questions
- Easy questions used to maintain flow
- Many question variants are possible
- Instructor must be prepared to use diverse questioning strategies

## Video (18 minutes)

- Excerpt from class taught at Southeastern Louisiana University in 1997
- Algebra-based general physics course
- *First Part:* Students respond to questions written on blackboard.
- *Second Part:* Students respond to questions printed in their workbook.

# Features of the Interactive Lecture

- High frequency of questioning
- Must often create unscripted questions
- Easy questions used to maintain flow
- Many question variants are possible
- Instructor must be prepared to use diverse questioning strategies

# High frequency of questioning

- Time per question can be as little as 15 seconds, as much as several minutes.
  - similar to rhythm of one-on-one tutoring
- Maintain small conceptual “step size” between questions for high-precision feedback on student understanding.

# Must often create unscripted questions

- Not possible to pre-determine all possible discussion paths
- Knowledge of probable conceptual sticking points is important
- Make use of standard question variants
- Write question and answer options on board  
*(but can delay writing answers, give time for thought)*

# Easy questions used to maintain flow

- Easy questions (> 90% correct responses) build confidence and encourage student participation.
- If discussion bogs down due to confusion, can jump start with easier questions.
- Goal is to maintain continuous and productive discussion with and among students.

# Many question variants are possible

- Minor alterations to question can generate provocative change in context.
  - add/subtract/change system elements (force, resistance, etc.)
- Use standard questioning paradigms:
  - greater than, less than, equal to
  - increase, decrease, remain the same
  - left, right, up, down, in, out

# Instructor must be prepared to use diverse questioning strategies

- If discussion dead-ends due to student confusion, might need to backtrack to material already covered.
- If one questioning sequence is not successful, an alternate sequence may be helpful.
- Instructor can solicit suggested answers from students and build discussion on those.



# Interactive Question Sequence

- Set of closely related questions addressing diverse aspects of single concept
- Progression from easy to hard questions
- Use multiple representations (diagrams, words, equations, graphs, etc.)
- Emphasis on qualitative, not quantitative questions, to reduce “equation-matching” behavior and promote deeper thinking

## Chapter 1 Electrical Forces


### “Flash-Card” Questions

#### In-Class Questions

##### Prerequisite Concepts:

- Positive and negative charges; Coulomb's law:  $F = kq_1q_2/r^2$
- Protons (+) and electrons (-)
- Superposition principle:  $F_{\text{net}} = F_1 + F_2 + \dots + F_n$
- Vector addition:  $F_{\text{net}x} = F_{1x} + F_{2x} + \dots + F_{nx}$
- Newton's second law,  $a = F/m$







Questions #1–2 refer to the figure below. Charge  $q_1$  is located at the origin, and charge  $q_2$  is located on the positive x axis, five meters from the origin. There are no other charges anywhere nearby.




1. If  $q_1$  is positive and  $q_2$  is negative, what is the direction of the electrical force on  $q_1$ ?
  - A. in the positive x direction
  - B. in the negative x direction
  - C. in the positive y direction
  - D. in the negative y direction
  - E. the force is not directed precisely along any of the coordinate axes, but at some angle
  - F. there is no force in this case
2. If  $q_1$  is positive and  $q_2$  is positive, what is the direction of the electrical force on  $q_1$ ?
  - A. in the positive x direction
  - B. in the negative x direction
  - C. in the positive y direction
  - D. in the negative y direction
  - E. the force is not directed precisely along any of the coordinate axes, but at some angle
  - F. there is no force in this case

3. In this figure, a proton is located at the origin, and an electron is located at the point (3m, 3m). What is the direction of the electrical force on the proton?


## “Flash-Card” Questions

- A. 
- B. 
- C. 
- D. 
- E. 
- F. 




4. In this figure, a proton is located at the origin, and a proton is located at the point (3m, 3m). The vector representing the electrical force on the proton *at the origin* makes what angle with respect to the positive x axis?

- A.  $0^\circ$   
 B.  $45^\circ$   
 C.  $90^\circ$   
 D.  $135^\circ$   
 E.  $225^\circ$   
 F.  $270^\circ$



5. In this figure, a proton is located at (0m, 5m) and an electron is located at the origin.



The electrical force on the electron is:


- A. directed toward the positive-y direction, and has greater magnitude than the electrical force acting on the proton
  - B. directed toward the positive-y direction, and has smaller magnitude than the electrical force acting on the proton
  - C. directed toward the positive-y direction, and has magnitude equal to the electrical force acting on the proton
  - D. directed toward the negative-y direction, and has greater magnitude than the electrical force acting on the proton
  - E. directed toward the negative-y direction, and has smaller magnitude than the electrical force acting on the proton
  - F. directed toward the negative-y direction, and has magnitude equal to the electrical force acting on the proton
6. Two charged particles are separated by a certain distance, and exert an electrical force on each other. What will happen to the magnitude of this electrical force if the separation between the particles is *decreased*?
- A. The force will decrease in magnitude.
  - B. The force will increase in magnitude.
  - C. The magnitude of the force will not change.
  - D. The magnitude of the force may decrease or increase, depending on whether the charges are like or unlike.
  - E. The magnitude of the force on one particle will increase, while that on the other particle will decrease.

7. Two particles with charges  $q_1$  and  $q_2$  are separated by a distance  $r$ . There are no other charges nearby. Consider the following actions:
- I. increase  $q_1$
  - II. increase  $q_2$
  - III. increase  $r$
  - IV. decrease  $r$


Which of the above actions will cause the magnitude of the force between the charges to *increase*?

- A. I and III only
  - B. I and IV only
  - C. II and III only
  - D. II and IV only
  - E. I and II and III
  - F. I and II and IV
8. When two charged particles are separated by 2 meters, the magnitude of the electrical force between them is  $F$ . What will be the magnitude of this force if their separation is increased to 4 meters?
- A.  $1/4 F$
  - B.  $1/2 F$
  - C.  $F$
  - D.  $2F$
  - E.  $4F$
  - F. not enough information given to determine magnitude of force
9. Which of these will result in the repulsive force between two identical charged particles *increasing* by a factor of 8:
- A. double one of the charges
  - B. double both of the charges
  - C. double one of the charges and cut the particle separation in half
  - D. triple one of the charges and cut the particle separation in half
  - E. triple both of the charges
  - F. double both of the charges and double the particle separation

10. A 6-C charge and a 12-C charge are separated by 2 m; there are no other charges present. Compared to the electrical force on the 6-C charge, the electrical force on the 12-C charge is:
- one-fourth as strong
  - one-half as strong
  - the same magnitude
  - two times as strong
  - four times as strong
11. In this figure, electrons are located on the y axis at  $y = 3$  m,  $y = 0$  m, and  $y = -3$  m. The direction of the net electrical force on the electron at the origin is:




- towards positive x
  - towards positive y
  - towards negative x
  - towards negative y
  - nowhere, since there is no net force on this electron
12. In this figure, protons are located on the x axis at  $x = -3$  m,  $x = 0$  m, and  $x = 1.5$  m. The direction of the net electrical force on the proton at the origin is:



- towards positive x
- towards positive y
- towards negative x
- towards negative y
- nowhere, since there is no net force on this proton


13. In this figure, positive charges of magnitude  $q$ ,  $q$ , and  $2q$  are located on the  $x$  axis as shown. The direction of the net electrical force on the positive charge at the origin is:

- A. towards positive  $x$   
 B. towards positive  $y$   
 C. towards negative  $x$   
 D. towards negative  $y$   
 E. nowhere, since there is no net force on this proton




14. In this figure, particles with charges  $-q$ ,  $+q$ , and  $-q$  are located on the  $x$  axis as shown. The direction of the net electrical force on the positive charge at the origin is:

- A. towards positive  $x$   
 B. towards positive  $y$   
 C. towards negative  $x$   
 D. towards negative  $y$   
 E. nowhere, since there is no net force on this proton



15. A electron is fixed at the origin; there are no other charges present. If a negative charge is brought in and released at a nearby point, and allowed to move freely, then as it moves the magnitude of the force acting on this negative charge will:
- A. always be zero  
 B. remain constant, but nonzero  
 C. always increase  
 D. always decrease, but never reach zero  
 E. sometimes increase and sometimes decrease  
 F. not enough information to decide

16. A electron is fixed at the origin; there are no other charges present. If a negative charge is brought in and released at a nearby point, and allowed to move freely, then as it moves the magnitude of the acceleration of this negative charge will:
- A. always be zero
  - B. remain constant, but nonzero
  - C. always increase
  - D. always decrease, but never reach zero
  - E. sometimes increase and sometimes decrease
  - F. not enough information to decide
17. A electron is fixed at the origin; there are no other charges present. If a negative charge is brought in and released at a nearby point, and allowed to move freely, then as it moves the speed of this negative charge will:
- A. always be zero
  - B. remain constant, but nonzero
  - C. always increase
  - D. always decrease, but never reach zero
  - E. sometimes increase and sometimes decrease
  - F. not enough information to decide
18. In this figure, a positive charge is located at the origin, and another positive charge is located at the point  $(-4\text{m}, 4\text{m})$ . The x component of the electrical force on the charge at the origin is:
- A. greater than zero
  - B. equal to zero
  - C. less than zero
  - D. may be equal to, less than, or greater than zero, depending on the precise magnitude of the two charges.






# Problem “Dissection” Technique

- Decompose complicated problem into conceptual elements
- Work through problem step by step, with continual feedback from and interaction with the students
- May be applied to both qualitative and quantitative problems

*Example: Electrostatic Forces*

**Figure 1. Diagram used for problem dissection**




Four charges are arranged on a rectangle as shown in Fig. 1. ( $q_1 = q_3 = +10.0 \mu\text{C}$  and  $q_2 = q_4 = -15.0 \mu\text{C}$ ;  $a = 30 \text{ cm}$  and  $b = 40 \text{ cm}$ .) Find the magnitude and direction of the resultant electrostatic force on  $q_1$ .

**Question #1:** How many forces (due to electrical interactions) are acting on charge  $q_1$ ?

(A) 0 (B) 1 (C) 2 (D) 3 (E) 4 (F) Not sure/don't know

**Figure 2. Direction options**



For questions #2-4 refer to Fig. 2 and pick a direction from the choices A, B, C, D, E, and F.

**Question #2:** Direction of force on  $q_1$  due to  $q_2$

**Question #3:** Direction of force on  $q_1$  due to  $q_3$

**Question #4:** Direction of force on  $q_1$  due to  $q_4$

Let  $F_2$ ,  $F_3$ , and  $F_4$  be the *magnitudes* of the force on  $q_1$  due to  $q_2$ , due to  $q_3$ , and due to  $q_4$  respectively.

**Question #5.**  $F_2$  is given by

- (A)  $kq_1q_2/a^2$
- (B)  $kq_1q_2/b^2$
- (C)  $kq_1q_2/(a^2 + b^2)$
- (D)  $kq_1q_2/\sqrt{a^2 + b^2}$
- (E) None of the above
- (F) Not sure/Don't know

**Question #6.**  $F_3$  is given by


- (A)  $kq_1q_3/a^2$
- (B)  $kq_1q_3/b^2$
- (C)  $kq_1q_3/(a^2 + b^2)$
- (D)  $kq_1q_3/\sqrt{a^2 + b^2}$
- (E) None of the above
- (F) Not sure/Don't know

(etc.)

# Workbook for Introductory Physics

Part II: Electricity and Magnetism, Optics, and Modern Physics

David E. Meltzer and Kandiah Manivannan



Iowa State University Physics Education Research Group